

T3/5/ALL

3/5/1

DIALOG(R)File 345:Inpadoc/Fam.& Legal Stat

(c) 2003 EPO. All rts. reserv.

16663698

Basic Patent (No,Kind,Date): GB 200031047 A0 20010131 <No. of Patents:
003>

PATENT FAMILY:

AUSTRALIA (AU)

Patent (No,Kind,Date): AU 200216204 A5 20020701

PERFUME COMPOSITIONS (English)

Patent Assignee: QUEST INTERNAT B V

Author (Inventor): BEHAN JOHN MARTIN; PERRING KEITH DOUGLAS; SMALL LES
; MCNULTY DAVID; RICHARDSON ANNE

Priority (No,Kind,Date): GB 200031047 A 20001220; WO 2001GB5589 W
20011218

Applic (No,Kind,Date): AU 200216204 A 20011218

IPC: * A61K-007/46; C11B-009/00; A61K-007/06; A61K-007/48; A61L-009/00
; C11D-003/50

Language of Document: English

GREAT BRITAIN (GB)

Patent (No,Kind,Date): GB 200031047 A0 20010131

PERFUME COMPOSITIONS (English)

Patent Assignee: QUEST INTERNAT B V

Priority (No,Kind,Date): GB 200031047 A 20001220

Applic (No,Kind,Date): GB 200031047 A 20001220

Language of Document: English

WORLD INTELLECTUAL PROPERTY ORGANIZATION, PCT (WO)

Patent (No,Kind,Date): WO 200249600 A1 20020627

PERFUME COMPOSITIONS (English)

Patent Assignee: QUEST INT (NL); BEHAN JOHN MARTIN (GB); PERRING
KEITH DOUGLAS (GB); SMALL LES (GB); MCNULTY DAVID (GB); RICHARDSON

ANNE (GB)

Author (Inventor): BEHAN JOHN MARTIN (GB); PERRING KEITH DOUGLAS
(GB); SMALL LES (GB); MCNULTY DAVID (GB); RICHARDSON ANNE (GB)

Priority (No,Kind,Date): GB 200031047 A 20001220

Applic (No,Kind,Date): WO 2001GB5589 A 20011218

Designated States: (National) AE; AG; AL; AM; AT; AU; AZ; BA; BB; BG;
BR; BY; BZ; CA; CH; CN; CO; CR; CU; CZ; DE; DK; DM; DZ; EC; EE; ES;
FI; GB; GD; GE; GH; GM; HR; HU; ID; IL; IN; IS; JP; KE; KG; KP; KR;
KZ; LC; LK; LR; LS; LT; LU; LV; MA; MD; MG; MK; MN; MW; MX; MZ; NO;
NZ; OM; PH; PL; PT; RO; RU; SD; SE; SG; SI; SK; SL; TJ; TM; TR; TT;
TZ; UA; UG; US; UZ; VN; YU; ZA; ZM; ZW (Regional) GH; GM; KE; LS;
MW; MZ; SD; SL; SZ; TZ; UG; ZM; ZW; AM; AZ; BY; KG; KZ; MD; RU; TJ;
TM; AT; BE; CH; CY; DE; DK; ES; FI; FR; GB; GR; IE; IT; LU; MC; NL;
PT; SE; TR; BF; BJ; CF; CG; CI; CM; GA; GN; GQ; GW; ML; MR; NE; SN;
TD; TG

Filing Details: WO 100000 With international search report

IPC: * A61K-007/46; C11B-009/00; A61K-007/06; A61K-007/48; A61L-009/00
; C11D-003/50

Derwent WPI Acc No: ; C 02-557636

Language of Document: English

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 June 2002 (27.06.2002)

PCT

(10) International Publication Number
WO 02/49600 A1

- (51) International Patent Classification⁷: **A61K 7/46**, C11B 9/00, A61K 7/06, 7/48, A61L 9/00, C11D 3/50
- (74) Agent: **KEITH W NASH & CO**; 90-92 Regent Street, Cambridge CB2 1DP (GB).
- (21) International Application Number: **PCT/GB01/05589**
- (22) International Filing Date:
18 December 2001 (18.12.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0031047.4 20 December 2000 (20.12.2000) GB
- (71) Applicant (*for all designated States except US*): **QUEST INTERNATIONAL B.V.** [NL/NL]; Huizerstraatweg 28, NL-1411 GP Naarden (NL).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BEHAN, John**, Martin [GB/GB]; The Forge, Boughton Aluph, Kent TN25 4JB (GB). **PERRING, Keith, Douglas** [GB/GB]; 14 Malvern Road, Ashford, Kent TN24 8HS (GB). **SMALL, Les** [GB/GB]; 20 Heathfield Road, Ashford, Kent TN24 8UD (GB). **MCNULTY, David** [GB/GB]; 9 Norfolk Street, Whitstable, Kent CT5 4HB (GB). **RICHARDSON, Anne** [GB/GB]; 6 Beech Avenue, Chartham, Canterbury, Kent CT4 7TA (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:**
— *of inventorship (Rule 4.17(iv)) for US only*
- Published:**
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/49600 A1

(54) Title: **PERFUME COMPOSITIONS**

(57) Abstract: A perfume composition comprises: e) at least about 25% by weight in total of at least five selected relaxing fragrance materials (R); f) optionally up to about 45% by weight in total of non-relaxing fragrance materials (NR), provided that the ratio by weight of R to NR is at least 0.75; g) optionally up to about 75% by weight in total of neutral fragrance materials (N); h) optionally up to about 25% by weight of other fragrance materials (M) provided that the weight ratio R to (M+NR) at least 0.75; and wherein all percentages are based on total weight of the fragrance materials consisting the perfume composition.

Title: Perfume Compositions**Field of the Invention**

This invention relates to perfume compositions which aim to promote relaxed and related mood states in subjects, particularly humans. The invention also concerns methods of delivering positive mood benefits or relaxation benefits to a subject.

Background to the Invention

It has long been known that fragrance materials and essential oils can promote feelings of relaxation and well-being. More recently these materials have been used in cosmetic products to similar effect *per se* and to enhance the effects of other materials/actives in the products. Many of these products rely on the use of simple mixes of essential oils to provide this benefit, eg lavender, jasmin oil, camomile and ylang-ylang. Many of the materials have been reported to promote different mood states, eg lavender has been reported to be both relaxing (see Torii et al in chapter 7, "Perfumery: the Psychology and Biology of Fragrances", S V Toller and G Dodd (eds), Chapman and Hall, 1991 reprint) and activating (see J S Jellinek, *Perfumer & Flavorist*, vol.22 (Sept/Oct), 29-41). It has therefore been difficult for formulators reliably to use literature information to make effective fragrances. The materials also are not optimised for their acceptability/hedonic effect but are often very basic odour combinations with little of the creativity found in more traditional fragrances.

Aromatherapy employs essential oils which are applied to the skin, for example with massage. The effect here involves additional processes other than smell. However, there are folk remedies in which essential oils are used primarily for their smell and a mood state change may result. In these examples the oils are used singly or in simple mixtures. There is an extensive literature concerning the activities and usage of essential oils for aromatherapy; see for example Tisserand ("The Art of Aromatherapy", C.W. Daniel, London (1985) and "Aromatherapy for Everyone", Penguin Books (1988) both by R. Tisserand). There is also a practice to use the term more freely, for example to claim "aromatherapy products" which may claim to "contain a fragrance which is relaxing and soothing". Consumers are aware that

aromatherapy products intend to enhance the sensory experience of the user, but there is no systematic teaching that can direct the perfumer to design products that would consistently provide this benefit. In many cases the claim is more akin to a marketing positioning statement, with a view to differentiating variants or competitive products.

Aroma-Chology® is a term coined by the Olfactory Research Fund Ltd. (see the extensive review by J Jellinek in *Cosmetics & Toiletries*, (1994) 109, pp 83 - 101). It is concerned specifically with the temporary, beneficial psychological effects of aromas on human behaviours and emotions to improve mood and quality of life. In fact, a large number of products promoted as having aromatherapy benefits can be more accurately identified for their Aroma-Chology® benefits as they produce temporary psychological effects. Again, however, there is no teaching as to how to formulate products to achieve such benefits qualitatively or quantitatively with a reliable expectation of success. In addition it is well known that fragrances can be perceived as associated with different attributes in different countries. Thus it may be that a fragrance found to be "stimulating" to the Japanese consumer will not be "stimulating" to the European consumer. Prior art perfumes do not exhibit the required robustness.

Hirsch (US Patent No.5904916) teaches a method for enhancing learning in a person by the administration of a mixed/floral odorant, preferably composed of fresh, citrus, herbaceous, fruity and floral odorants, exemplified by Mixed-Floral IFF No. 2635-AS. No further teaching was disclosed on other fragrances of the invention which may have similar psychological benefits.

Clow and Huckelbridge have shown that the smell of chocolate can modulate levels of secretory immunoglobulin A (IgA) which has been linked to stress and mood (see A A Stone et al in *J. Personality and Social Psychology*, (1987), volume 52(5), 988-993). No teachings were provided as to how this might be related to perfume development.

Japanese Patent No.9-227399 (1997) claims extracts of plants of the Labiatae genus.

Research on the neural basis of emotion has concentrated mostly on fear. The amygdala has been identified as a key component of a network of neural pathways in fear conditioning (see J Le Doux, Annual Review of Psychology, (1995), volume v46, 209) and emotional memories. Many human emotions exist as part of complex neural systems that have evolved to help us survive. Le Doux ("The Emotional Brain: The mysterious Underpinnings of Emotional Life", (1996), publ. Simon & Schuster Inc, New York) points out that emotional responses are hard-wired into the brain's circuitry, and that the stimuli that trigger emotional states are learned through experience. These can be uniformly perceived throughout a population, for example, the sight of a snake triggering a fear response, or of a picture of an injured child eliciting a depressed response. In contrast, the invention described herein identifies a class of perfumes which elicit similarly uniform but pleasant responses. Other fragrances are more susceptible to interpretation being governed by context, eg the country in which the assessor grew up, the appearance of the product in which the perfume is assessed, the use of the product (eg washing with soap).

Kan *et al* (18th International Congress, IFSCC, Venezia, (Oct 1994), preprints 769-784) showed that cosmetics which beautify the person have psychoimmunological effects which manifest as an increase in the body's immune status, as measured by salivary IgA. Japanese Patent No. 06172781 describes deactivating (sedative) perfumes based on the use of 1,3-dimethoxybenzene-5-methylbenzene. No teaching on other perfume ingredients was disclosed in either publication.

Alaoui-Ismaili *et al* (Chem.Senses (1997), 22, 237-248) sought evidence of a linkage between emotion and the sense of smell, particularly smell preference. The results did not provide evidence for the postulated preferential link between olfaction and emotion (although preference scores for their odorants agreed with literature values), leading to the conclusion that the emotional content of smell cannot be predicted from preference. Van Toller *et al* (Chemical Senses, (1993) vol.18, pp 1-16) found that increased electrical activity (alpha waves) in the brain was associated with increased pleasantness, intensity and familiarity for a range of widely disparate odours.

The present invention seeks to address at least some of the problems described above, in particular that of the creative scope available to perfumers, and possibly also robustness across different consumer groups and performance reliability.

Summary of Invention

The present invention relates to perfume compositions which aim to induce or be associated with positive, low activation moods and emotions. In a first aspect of the invention, a perfume composition comprises:

- a) from about 25% in total of at least five Relaxing Ingredients (R);
- b) optionally up to 45%, preferably up to 35%, more preferably up to 25%, in total of Non-Relaxing Ingredients (NR), provided that the ratio of R to NR is at least 0.75, preferably at least 0.9;
- c) optionally up to 75% in total of Neutral Ingredients (N);
- d) optionally up to 25% of other perfumery materials (M) provided that the weight ratio R to (M+NR) exceeds 0.75, preferably 0.9;

and wherein (i) all percentages are based on weight of the perfume composition;

(ii) 'R' ingredients comprise anethole, Bangalol™, basil oil, cis-hex-3-enol, coumarin, ethylene brassylate, ethyl linalol, Florosa™, Galaxolide™, geraniol, cyclohexadecanolide, cyclopentadecanone, methyl anthranilate, alpha-iso-methyl ionone, Prunella™, Silvanone™, alpha-terpineol, Traseolide™, Ultravani™, gamma-undecalactone, vetiver oil, vetiver acetate;

(iii) 'NR' ingredients comprise methyl nonyl aldehyde (aldehyde MNA), allyl amyl glycolate, acetyl cedrene, Amberlyn Super™, amyl salicylate, armoise oil, benzyl salicylate, bergamot oil, Bourgeonal™, cedar leaf oil, citronellol, beta-damascone, dimethyl benzyl carbonyl acetate, Ethyl Safranate™, Everniate™, geranyl nitrile, Helional™, heliotropin, hexyl salicylate, lemon oil, Ligustral™, Lillal™, Lyrall™, Mefrosol™, orange oil, orange terpenes, tagetes oil, tetrahydrogeraniol, vanillin;

(iv) 'N' ingredients comprise benzyl acetate, cassis base, Cyclamen Aldehyde™, carvone, cinnamic alcohol, dihydroeugenol,

dihydromyrcenol, eugenol, Extralide™, galbanum, gamma-decalactone, hydroxycitronellal, indole, isoeugenol, jasmin oil, Jasmopyrane Forte™, linalol, linalyl acetate, methyl dihydrojasmonate (MDJ), octahydrocoumarin, patchouli oil, 2-phenylethyl alcohol, rose oxide, rose oil, Sandalone™, Sandalore™, styrallyl acetate, ylang-ylang;

(v) 'M' ingredients comprise perfumery materials not included in the above, excluding odourless or low-odour solvents or diluents used as vehicles for other perfume components.

Another aspect of the invention relates to a method for delivering positive mood benefits, particularly relaxation, to a subject, particularly a human subject, comprising delivering a perfume composition in accordance with the invention in a form selected from consumer products intended for application to skin, hair, hard surface or fabrics, and from air care products such as airfresheners.

The present invention is based on extensive testing of fragrance materials, both by consumer testing and by measurement of brain electrical activity particularly alpha wave activity measured by electroencephalography (EEG), and statistical analysis of the resulting data to classify the materials into different categories, namely relaxing fragrance materials or ingredients (R) that induce in subjects exposed to them positive, low activation moods and emotions, such as relaxation (i.e. relaxing properties), non-relaxing fragrance materials or ingredients (NR) that induce in subjects exposed to them negative, high activation moods or emotions (i.e. non-relaxing properties), neutral fragrance materials or ingredients (N) having a neutral effect in terms of relaxing properties. Other fragrance materials (of which there are around three to four thousand currently available commercially and used in perfume formulation) are designated as class M materials. Based on this classification of fragrance materials, the invention enables perfume compositions to be defined that are likely to induce in subjects exposed to them positive, low activation moods and emotions, such as relaxation. Such perfume compositions are referred to herein for convenience as relaxing perfumes. The definition of the perfume composition nevertheless provides sufficient freedom in formulation to permit consideration of the hedonic properties of the composition. The invention can thus

enable formulation of perfume compositions that are relaxing and also have good hedonic properties.

The present invention describes how to formulate reliably novel fragrances which are likely to induce or be associated with positive, low activation moods and emotions, particularly relaxing effects. The effects are sufficiently pronounced that they can be measured reliably and reproducibly. The perfume compositions made according to the teachings disclosed herein can be hedonically pleasant, suitable for a wide range of consumer products, and of sufficient pleasantness/acceptability that they would be appropriate even if they did not possess added functionality. In addition, perfume compositions of the invention can be resilient to variation in the target consumer group (eg Japanese versus American) and have been found to be perceived as consistently relaxing/reassuring etc for consumers in England, France, USA and Japan.

Perfume compositions in accordance with the invention have been found:

- a) to promote a reduction in the amplitude of alpha wave activity in the brain;
- b) to promote positive mood states such as relaxation. In tests subjects have reported that they feel more relaxed after smelling or using consumer products incorporating the perfume compositions, and subjects report that the products themselves smell more relaxing;
- c) to promote calming, warming, sensual, caring, reassuring, safe mood states;
- d) not to promote negative mood states such as depressing, stressful, irritating, or bored mood states.

The prior art shows that fragrances which are relaxing (or promote related affect responses) operate by mechanisms other than pleasantness alone. Mood and emotion are hard-wired in the brain and are triggered by external stimuli. The basis for systematically selecting olfactory stimuli which evoke specific patterns of brain activity and evoke specific emotions were not previously known. There are a number of methods for gauging the state of relaxation, for example self-appraisal based on adjective-lists, and measurements of brain wave activity, particularly that of the alpha waves (8 to 13Hz range). In other sensory modalities it would be expected that an increase in power/amplitude in the alpha frequency band would be associated with positive feelings. However, we have found unexpectedly that reduction of odour-derived

alpha activity is associated with increased positive feelings and in particular we have found that it is related to positive feelings of less activated mood states (see previous reference for description of positive/negative and activated/deactivated moods and emotions). We have in particular found that reported relaxation is enhanced by fragrances associated with reduced alpha wave activity. We were able to identify fragrance ingredients which are used at higher levels in relaxing fragrances. These were generally materials with notes such as sweet, musky, fruity, floral and amber i.e. unlike the materials disclosed in US 5904916 and JP 9-227399 referred to above. Increasing the level of these relaxing ingredients (referred to as class 'R' here) increased the likelihood that the fragrance would have a suitable character to deliver the relaxing benefit. Other ingredients reduce the likelihood that the benefit will be achieved as their level in the fragrance is increased, ie they were the non-relaxing group (referred to as class 'NR' here). A third group had a neutral effect (class 'N'). Combinations of fragrance ingredients which are commercially useful and aesthetically pleasant generally require materials in all three classes to be present. The number of perfume ingredients available commercially is around three to four thousand materials. Surprisingly, these other materials (not in class 'R', 'NR' or 'N' and designated herein as class 'M') may be added to fragrances of the invention without loss of benefit, provided that the weight ratio of relaxing materials to the sum of class M and class NR materials is equal to or greater than around 0.75. In the perfume art, some materials having no odour or very weak odours are used as diluents or vehicles for other ingredients. Non-limiting examples of these are dipropylene glycol (DPG), diethyl phthalate, benzyl benzoate, triethyl citrate and isopropyl myristate. Such materials are not counted towards the definition/formulation of the perfume compositions of the invention.

The present invention thus provides a perfume composition comprising:

- (a) at least about 25% by weight in total of at least five relaxing fragrance materials (R);
- (b) optionally up to about 45% by weight in total of non-relaxing fragrance materials (NR), provided that the weight ratio of R to NR is at least 0.75;
- (c) optionally up to about 75% by weight in total of neutral fragrance materials (N);
- (d) optionally up to about 25% by weight of other fragrance materials (M) provided that the weight ratio R to (M+NR) is at least 0.75;

- and wherein
- (i) all percentages are based on total weight of the fragrance materials consisting the perfume composition excluding any solvents and diluents;
 - (ii) the R materials are selected from the group consisting of anethole, 2-ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)but-2-enol (eg Bangalol™), basil oil, cis-hex-3-enol, coumarin, ethylene brassylate, ethyl linalol, 2-(2'-methylpropyl)-4-hydroxy-4-methyltetrahydropyran (eg Florosa™), hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-benzopyran (eg Galaxolide™), geraniol, cyclohexadecanolide, cyclopentadecanone, methyl anthranilate, alpha-iso-methyl ionone, mixtures of dimethyl benzyl carbonyl butyrate and phenoxyethyl isobutyrate (e.g. Prunella™), mixtures of cyclohexadecanolide and cyclopentadecanone (e.g. Silvanone™), alpha-terpineol, 6-acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin (eg Traseolide™), 2-ethoxy-4-methylphenol (eg Ultravani™), gamma-undecalactone, vetiver oil, vetiver acetate;
 - (iii) the NR materials are selected from the group consisting of methyl nonyl aldehyde, allyl amyl glycolate, acetyl cedrene, 3a,6,6,9a-perhydrotetramethylnaphtho [2,1-b] furan (eg Amberlyn Super™), amyl salicylate, armoise oil, benzyl salicylate, bergamot oil, 4-tert-butyl-3-phenylpropanal (eg Bourgeonal™), cedar leaf oil, citronellol, beta-damascone, dimethyl benzyl carbonyl acetate, ethyl 2,6,6-trimethylcyclohexadienecarboxylate (eg Ethyl Safranate™), methyl 2,4-dihydroxy-3,6-dimethylbenzoate (eg Everniate™), geranyl nitrile, 3-(1,3-benzodioxol-5-yl)-2-methylpropanal (eg Helional™), heliotropin, hexyl salicylate, lemon oil, 2,4-dimethyl-3-cyclohexene-1-carbaldehyde (eg Ligustral™), 3-(4-(1,1-dimethylethyl)phenyl)-2-methylpropanal (eg Lilial™), 4-(4-hydroxy-4-methylpentyl)cyclohex-3-enecarbaldehyde (eg Lyrall™), 3-methyl-5-phenylpentanol (eg Mefrosol™), orange oil, orange terpenes, tagetes oil, tetrahydrogeraniol, vanillin;
 - (iv) the N materials are selected from the group consisting of benzyl acetate, cassis base, 2-methyl-3-(4-(1-methylethyl)phenyl)propanal (eg Cyclamen Aldehyde™), carvone, cinnamic alcohol, dihydroeugenol, dihydromyrcenol, eugenol, 6-acetyl-1,1,3,4,4,6-

hexamethyltetrahydronaphthalene (eg Extralide™), galbanum, gamma-decalactone, hydroxycitronellal, indole, isoeugenol, jasmin oil, 3-pentyltetrahydro-2H-4-pyranyl ethanoate (eg Jasmopyrane Forte™), linalol, linalyl acetate, methyl dihydrojasmonate, octahydrocoumarin, patchouli oil, 2-phenylethyl alcohol, rose oxide, rose oil, isobornylcyclohexanol (eg Sandalone™), 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol (eg Sandalore™), styrallyl acetate, ylang-ylang;

(v) the M materials comprise perfumery materials not included in the above, excluding odourless or low-odour solvents or diluents used as vehicles for fragrance materials.

Preferred perfume compositions comprise at least 35% by weight in total of class R materials, even more preferably at least 45% by weight in total of class R materials.. Also preferred are those perfumes wherein the weight ratio of R to NR (or R to the sum of NR and M) exceeds unity, or more preferably exceeds two, or is even higher

The perfume composition preferably includes optionally up to about 25% by weight in total of non-relaxing (NR) fragrance materials.

In a particularly preferred embodiment at least 5% by weight, or even more preferably at least 10% by weight, of the perfume composition comprises class R ingredients drawn from the following list, mostly characterised by exhibiting sweet and/or musky notes: coumarin, ethylene brassylate, Galaxolide™, cyclohexadecanolide, cyclopentadecanone, Traseolide™, Ultravanil™, gamma-undecalactone.

The perfume composition may optionally include one or more odourless or low-odour solvents and/or diluents, e.g. as a vehicle for a fragrance material. Any such solvents and/or diluents are not included when calculating percentages and ratios of R, NR, N and M materials of the composition.

It is possible to combine preferred embodiments as defined above to produce fragrances which are highly preferred.

Proprietary Fragrance Materials

Further details on materials are given below.

Name	Supplier	Technical name/Alternative names *
Anther™	Q	Phenylether isoamyl ether
Amberlyn Super™	Q	3a,6,6,9a-perhydrotetramethylnaphtho [2,1-b] furan / Ambrox™ / Cetalox™
Bangalol™	Q	2-Ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)but-2-enol
Beauvertate™	Q	Methyl 2-nonenoate
Bourgeonal™	Q	4-tert-butyl-3-phenylpropanal
Cyclamen aldehyde™G-R		2-methyl-3-(4-(1-methylethyl)phenyl)propanal
Ethyl saframate™	Q	Ethyl 2,6,6-trimethylcyclohexadienecarboxylate
Everniate™	Q	Methyl 2,4-dihydroxy-3,6-dimethylbenzoate / Oakmoss synthetic
Extralide™	Q	6-Acetyl-1,1,3,4,4,6-hexamethyltetrahydronaphthalene / Tonalid™
Florosa™	Q	2-(2'-methylpropyl)-4-hydroxy-4-methyltetrahydropyran
Galaxolide™	IFF	Hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-Benzopyran
Indolal	D	4,5-(1,2-indano)-1,3-dioxan
Helional		3-(1,3-benzodioxol-5-yl)-2-methylpropanal
Heliotropin		1,3-benzodioxole-5-carbaldehyde / Piperonal
Iso Ambois™	Q	1 - (1,2,3,4,5,6,7,8 - Octahydro - 2,3,8,8 -tetramethyl - 2 - naphthyl)ethan - 1 - one / Iso E Super
Jasmpyrane Forte™	Q	3-pentyltetrahydro-2H-4-pyranyl ethanoate
Ligustral™	Q	cis and trans 2,4-dimethyl-3-cyclohexene-1-carbaldehyde
Lilial™	G-R	3-(4-(1,1-dimethylethyl)phenyl)-2-methylpropanal

Lyrall™	IFF	4-(4-hydroxy-4-methylpentyl)cyclohex-3-enecarbaldehyde
Mefrosol™	Q	3-methyl-5-phenylpentanol
Melonal™	G	2,6-Dimethyl-5-heptenal
Ozonal base™	Q	complex composition
Prunella™	F	a mixture of dimethyl benzyl carbinyl butyrate and phenoxyethyl isobutyrate
Sandalone™	Q	isobornylcyclohexanol
Sandalore™	G-R	5-(2,2,3-Trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol
Silvanone™	Q	Cyclopentadecanone/Cyclohexadecanolide
Traseolide™	Q	6-Acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin
Ultravani™	Q	2-Ethoxy-4-methylphenol
Vetiver acetate	IFF	1,2,3,3a,4,5,6,8a-octahydro-2-isopropylidene-4,8-dimethyl azulen-6-yl acetate

* Based on major species present in the component.

Suppliers:

D:	Dragoco
F:	Firmenich
G-R	Givaudan-Roure
IFF:	International Flavors and Fragrances
Q:	Quest International

Materials of class 'M' include prior art perfume materials which are not specified as being members of any of classes R, NR or N, excluding odourless or low-odour solvents or diluents, as noted above. They may be single ingredients, or mixtures both synthetic and natural (for example essential oils, and concretes), and are well described e.g. in: "Common Fragrance and Flavor Materials" by Bauer, Garbe and Surburg, VCH Publ., 2nd edition (1990), and "Perfume and Flavour Materials", Steffen Arctander, published in two volumes by the author (1969), and also by Arctander "Perfume and Flavor Materials of Natural Origin", (1960).

Another aspect of the invention relates to a method for delivering positive mood benefits; particularly relaxation, to subjects, particularly humans, comprising delivering fragrance in a

form selected from consumer products intended for application to the body (that is to skin or hair), to hard surfaces (eg kitchen and bathroom worktops, ceramic surfaces), to fabrics, and for air care benefits (for example, airfresheners). Such products can take a variety of forms including powders, bars, sticks, tablets, creams, mousses, gels, liquids, sprays, and sheets. The amount of perfume in such products may lie in a range from 0.05% (as for example in low odour skin creams) to 30% (as for example in fine fragrances) by weight thereof. The incorporation of perfume into products of these types is known, and existing techniques may be used for incorporating perfumes for this invention. It may be possible to incorporate perfume directly into a product, but another possibility is, to absorb the perfume on a carrier material and then admix the perfume-plus-carrier mixture into the product.

The invention thus also includes within its scope a consumer product, e.g. a fabric treatment composition, a personal product composition or a hard surface treatment composition, comprising a perfume composition in accordance with the invention.

Also included within the scope of the invention is a method of delivering positive mood benefits or relaxation benefits to a subject, particularly a human, comprising administering to the subject an effective amount of a perfume composition in accordance with the invention. The composition should be administered in appropriate amount to produce a benefit (i.e. a suprathreshold amount) without causing irritation (i.e. a non-irritant amount). An appropriate effective amount of any given composition can be readily determined e.g. by experiment. To be effective, the composition should be administered for inhalation by the subject.

The invention will be further described, by way of illustration, in the following, non-limiting examples. Reference is made to the accompanying drawings, in which:

Figure 1 is a schematic diagram illustrating electrode placement on the head of a test subject; and

Figure 2 is an example of a map obtained by multidimensional scaling methods (MDS) analysis of brain electrical activity mapping (BEAM) data.

In the Examples, all amounts are expressed as percentages by weight, unless otherwise specified.

EXAMPLES 1-6: Relaxing perfume compositions incorporating class NR materials.

COMPOSITIONS

w/w%

INGREDIENT	Class	R1	R2	R3	R4	R5	R6
ACETYL CEDRENE	NR	4.32	8.57	3.09	10.41	2.91	4.13
ALDEHYDE MNA	NR	0.04	0.02	0.03	0.04	0.03	0.03
ALLYL AMYL GLYCOLATE	NR	0.16	0.21	0.26	0.24	0.15	0.16
AMBERLYN SUPER (Q)	NR	0.13	0.17	0.13	0.20	0.13	0.13
AMYL SALICYLATE	NR	2.53	2.20	1.88	1.76	2.44	2.43
BANGALOL (Q)	R	1.15	0.73	0.42	0.64	0.43	0.76
BEAUVERTATE (Q)	M	0.00	0.00	0.00	0.44	0.00	0.00
BENZYL ACETATE	N	0.92	3.75	3.06	2.44	1.91	1.67
BENZYL SALICYLATE	NR	2.65	4.60	6.04	7.68	6.61	5.95
BOURGEONAL	NR	0.14	0.44	0.25	0.44	0.17	0.27
CARVONE	N	0.11	0.10	0.10	0.14	0.10	0.11
CINNAMIC ALCOHOL	N	0.72	0.33	0.35	0.37	0.52	0.64
CITRONELLOL	NR	1.04	1.53	0.93	1.41	1.65	1.35
DIHYDRO MYRCENOL	N	2.42	1.02	1.95	1.89	2.25	2.37
ETHYLENE BRASSYLATE	R	1.31	1.70	1.19	1.45	1.13	1.26
EXTRALIDE(Q)	N	0.21	0.00	0.00	0.00	0.00	0.00
FLOROSA (Q)	R	9.32	10.51	11.85	9.43	2.90	8.89
GALBANUM	N	0.00	0.00	0.20	0.00	0.00	0.00
HEXYL SALICYLATE	NR	1.30	5.80	6.50	5.03	3.84	3.08
ISO AMBOIS (Q)	M	0.00	0.06	0.00	0.00	0.00	0.00
JASMIN	N	0.33	0.87	0.68	0.88	0.27	0.53
JASMOPYRANE FORTE (Q)	N	2.50	2.47	1.51	1.87	2.30	2.33
LIGUSTRAL (Q)	NR	0.12	0.25	0.09	0.15	0.15	0.13
LINALOL	N	2.56	1.32	6.97	3.41	1.35	2.48
LINALYL ACETATE	N	3.39	4.71	4.14	5.20	2.92	3.26
LYRAL (IFF)	NR	4.18	3.19	3.72	6.40	3.73	3.78
MDJ	N	6.66	10.95	10.24	3.97	1.67	7.48
MEFROSOL (Q)	NR	2.93	5.38	7.71	4.92	4.89	3.51
METHYL IONONE ALPHA ISO	R	19.26	6.15	4.84	6.32	1.85	12.50
OCTAHYDROCOUMARIN	N	0.00	0.00	0.00	0.00	0.19	0.00
ORANGE OIL	NR	1.40	2.76	2.47	2.60	1.46	1.46
PHENYL ETHYL ALCOHOL	N	0.94	6.75	3.21	2.78	1.69	1.80
SANDALONE(Q)	N	2.81	2.98	2.13	2.39	2.64	2.66
SILVANONE (Q)	R	1.47	2.11	1.96	1.76	2.05	1.70
TERPINEOL ALPHA	R	1.45	2.16	1.88	2.07	1.67	1.62

TRASEOLIDE (Q)	R	21.26	5.95	10.09	11.08	43.83	21.11
TUBEROSE BASE	M	0.00	0.00	0.00	0.00	0.00	0.21
ULTRAVANIL (Q)	R	0.01	0.01	0.00	0.00	0.01	0.01
UNDECALACTONE GAMMA	R	0.26	0.25	0.13	0.19	0.16	0.20
total:		100.00	100.00	100.00	100.00	100.00	100.00
sum R		55.49	29.57	32.35	32.94	54.01	48.07
sum NR		20.94	35.13	33.09	41.28	28.18	26.40
sum N		23.57	35.25	34.55	25.34	17.82	25.33
sum M		0.00	0.06	0.00	0.44	0.00	0.21
Ratio R/NR		2.65	0.84	0.98	0.80	1.92	1.82
Ratio R/(M +NR)		2.65	0.84	0.98	0.79	1.92	1.81

Notes: a) Materials marked 'Q' are trademark materials available from Quest International

b) Materials marked 'IFF' are trademark materials available from International Flavors and Fragrances

EXAMPLES 7-8: Comparative examples of relaxing (R7) and non-relaxing (NR1) perfume compositions.

INGREDIENT	w/w%		
	Class	R7	NR1
ACETYL CEDRENE	NR	0.00	14.58
ALDEHYDE MNA	NR	0.00	0.05
ALLYL AMYL GLYCOLATE (Q)	NR	0.00	0.36
AMBERLYN SUPER PM (Q)	NR	0.00	0.27
BANGALOL (Q)	R	1.46	0.00
BENZYL ACETATE (Q)	N	5.48	6.84
CARVONE LAEVO (Q)	N	0.15	0.18
DIHYDRO MYRCENOL (Q)	N	4.38	5.47
ETHYLENE BRASSYLATE	R	1.68	0.00
FLOROSA	R	13.14	0.00
HYDROXYCITRONELLAL	N	7.30	9.11
JASMIN	N	0.95	1.18

JASMOPYRANE FORTE (Q)	N	3.94	4.92
LIGUSTRAL (Q)	NR	0.00	0.55
LINALOL	N	7.56	9.43
LINALYL ACETATE	N	5.84	7.29
LYRAL	NR	0.00	6.84
MDJ	N	19.43	24.27
METHYL IONONE ALPHA ISO	R	18.25	0.00
ORANGE OIL	NR	0.00	2.28
PHENYL ETHYL ALCOHOL	N	5.11	6.38
SILVANONE (Q)	R	2.92	0.00
TERPINEOL ALPHA	R	2.19	0.00
UNDECALACTONE GAMMA	R	0.22	0.00
total:		100%	100%
Sum R		39.86	0.00
Sum NR		0.00	24.93
Sum N		60.14	75.07
Sum M		-	-
Ratio R/NR		-	-
Ratio R/(M+NR)		-	-

EXAMPLE 9: Relaxing perfume composition (R8) incorporating significant proportion of class M materials

Ingredient	Class	w/w% R8
ACETYL CEDRENE	NR	1
ANTHER (Q)	M	0.4
BANGALOL (Q)	R	0.4
BEAUVERTATE (Q)	M	0.3
BENZYL ACETATE	N	2
BOURGEONAL(Q)	NR	0.6
CEDAR LEAF OIL (10% in DPG)	NR	0.6
ETHYL LINALOL	R	9
ETHYL LINALYL ACETATE	M	2
ETHYL SAFRANATE (Q) (10% in DPG)	NR	0.2
FLOROSA (Q)	R	18
GERANYL NITRILE (10% in DPG)	NR	0.9

HEXYL CINNAMIC ALDEHYDE	M	9
INDOLAL (10%) DPG	M	0.1
JASMOPYRANE FORTE (Q)	N	1
MEFROSOL (Q)	NR	4.5
MENTHANYL ACETATE	M	3
METHYL IONONE ALPHA ISO	R	4
METHYL ISO EUGENOL	M	0.3
MOSS OAKMOSS SYNTHETIC (10% in DPG)	NR	0.2
OZONAL BASE (Q)	M	0.3
PATCHOULI OIL ACID WASHED	N	1
PHENYL ETHYL ALCOHOL	N	4
PARTA-TERT.BUTYL CYCLOHEXYL ACETATE	M	3
SILVANONE (Q)	R	4
TERPINEOL ALPHA	R	3
TETRAHYDROGERANIOL	NR	1
TRASEOLIDE (Q)	R	25
ULTRAVANIL (Q) (1% in DPG)	R	0.7
YLANG-YLANG	N	0.5
	total	100
class analysis (excluding solvent)	sum R	65.22
	sum NR	7.42
	sum N	18.64
	sum M	18.72
	Ratio:	
	R/NR	8.79
	Ratio:	7.27
	R/(M+NR)	2.50

EXAMPLE 10: Further embodiments of perfume compositions (R9 to R12) in accordance with the invention, with good hedonic properties.

INGREDIENT	Class	w/w%			
		R9	R10	R11	R12
ACETYL CEDRENE	NR	4	5	3.5	0
ANETHOLE	R	0.1	0	0	0
ANISIC ALDEHYDE	M	0	0	0.5	1
ANTHER	M	0.8	0	0	0
BANGALOL (Q)	R	0.5	0	0.5	1.4
BENZYL ACETATE	N	4	0	0	1
BOURGEONAL (Q)	NR	0.3	0.5	0.2	0.1

CINNAMIC ALCOHOL	N	0	0	0	0.8
CIS 3 HEXENOL	R	0.2	0	0	0
CIS 3 HEXENYL ACETATE	M	0.2	0	0	0
CITRONELLOL	NR	0	0	2	1.5
DECALACTONE GAMMA	N	0	0.3	0	0
DEWFRUIT BASE (Q)	M	0.1	0	0	0
DIHYDROEUGENOL	N	0	0.3	0	0.1
DIPROPYLENE GLYCOL	Solv	6.6	25.6	10	9.6
ETHYL LINALOL	R	10	4	1.5	0
ETHYL LINALYL ACETATE	M	5	0	3.5	1
ETHYL SAFRANATE (10% in DPG)	NR/Solv	0	0.2	0	0.6
ETHYL VANILLIN	M	0	0	0	0.5
FLOROSA (Q)	R	15	15.2	3	10
GERANYL NITRILE (10% in DPG)	NR/Solv	2	0.2	0	0
HELIOTROPIN	NR	0	0	0	0.9
HEXYL CINNAMIC ALDEHYDE	M	0	5	2	0
HEXYL SALICYLATE	NR	9	0	0	1
INDOLAL (10% in DPG)	M/Solv	0.3	0.9	0	0
IONONE	M	0	1.5	0	0
ISO AMBOIS SUPER (Q)	M	0	0	0	15
JASMATONE	M	0.1	0	0	0
JASMOPYRANE FORTE (Q)	N	2	0	0	0
LIGUSTRAL (Q)	NR	0.1	0	0	0
MEFROSOL (Q)	NR	10	3	6	3
MELONAL (10% in DPG)	M	0	0.3	0	0
METHYL ANTHRANILATE (10% in DPG)	R/Solv	0.3	0	0	0
METHYL IONONE, alpha iso-	R	0	0	2	25
MOSS OAKMOSS (10% in DPG)	M	4	0	0	0
OCTAHYDROCOUMARIN	N	0	0	1	0
ORANGE TERPENES	NR	4	0	0	0
OZONAL BASE	M	0.5	6	0	0
PHENYL ETHYL ALCOHOL	N	6	0	2	0
PRUNELLA	R	0	0	0	2
SILVANONE (Q)	R	0	2	2.5	1.5
TETRAHYDROGERANIOL	NR	1.5	3	2.5	0
TRASEOLIDE (Q)	R	12	27	55	23
ULTRAVANIL (1% in DPG) (Q)	R/Solv	0	0	1	1
UNDECALACTONE GAMMA	R	0.2	0	0	0
YLANG YLANG	N	1.2	0	1.3	0
TOTAL		100	100	100	100
Class Analysis (excl.solvent)					
R		41.76	65.82	72.48	70.79
NR		31.96	15.76	15.95	7.38
N		14.50	0.82	4.83	2.14

M	11.78	17.60	6.74	19.69
R/NR	1.31	4.18	4.54	9.59
R/(NR+M)	0.95	1.97	3.19	2.61

Odour descriptions of the perfume compositions listed above are as follows:

<u>Composition</u>	<u>Odour Type</u>
R1	Floral fruity
R2	Muguet woody amber
R3	Green fruity floral
R4	Green floral
R5	Powdery floral
R6	Fruity floral
R7	Muguet violet sandalwood
R8	Fruity floral
R9	Green fruity floral
R10	Marine floral
R11	Powdery floral
R12	Floral fruity

EXAMPLES 11-12 Mood data

Example 11.

Methodology

The moods elicited by different fragrances were investigated using naïve consumers. Each subject assessed each of the test fragrances, presented blind and in a balance randomised order. The method used was line-scaling. In this method a mark is placed on a line at a point between the two ends which represents the relative similarity/difference between the properties of the sample and the adjectives associated with each end of the line. The instructions given were "Please put a mark on each line listed to indicate how you feel about the odour". This

was followed by a list of the attributes, shown below, each with a line scale as illustrated. The data were analysed using standard statistical methods, see below.

The characteristics assessed were as follows:

Trendy	Calming
Nostalgic	Relaxing
Warm	Comforting
Simulating	Happy
Modern	Irritating
Sensual	Depressing
Stressful	Invigorating
Soothing	Cooling
Refreshing	Sexy
Safe	Reassuring
Caring	Liking

Example of line scale

CALMING

ExtremelyX..... Not at all

The data was analysed by Analysis of Variance (ANOVA) to generate mean scores which were used for further analysis.

<u>Fragrance</u>	<u>Sum of 'Relaxation' scores**</u>	<u>Effect</u>
R1	80	Highly Relaxing
R2	62	Relaxing
R3	59	Relaxing
R4	53	Relaxing

R5	77	Highly Relaxing
R6	66	Relaxing
R7	85	Highly Relaxing
R8	75	Highly Relaxing
NR1	25	Non-Relaxing
2635-AS*	50	Relaxing

*Prior art fragrance in US 5904916

**estimates

Example 12

Methodology

The moods elicited by different fragrances were investigated using naïve consumers. Each subject assessed each of the test fragrances, presented blind and in a balance randomised order. The method used was line-scaling. In this method a mark is placed on a line at a point between the two ends which represents the relative similarity/difference between the properties of the sample and the adjectives associated with each end of the line. The instructions given were "Please put a mark on each line listed to indicate how the product makes you feel". The characteristics used were as follows:

Trendy	Calm
Nostalgic	Warm
Stimulated	Modern
Sensual	Stressed
Soothed	Refreshed
Safe	Cared for
Relaxed	Comfortable
Happy	Irritated
Depressed	Invigorated
Sexy	Reassured

Note: descriptors should be understandable and relevant to local tastes and understanding.
Words such as 'cool; can be ambiguous when used internationally.

<u>Fragrance</u>	<u>Sum of 'Relaxation' scores**</u>	<u>Effect</u>
R1	75	Highly Relaxing
R2	66	Relaxing
R3	61	Relaxing
R4	50	Relaxing
R5	71	Highly Relaxing
R6	67	Relaxing
R7	78	Highly Relaxing
R8	71	Highly Relaxing
NR1	29	Non- Relaxing
2635-AS*	46	Relaxing

*Prior art fragrance in US 5904916

** estimates

Example 13: Spontaneous cortical activity in response to a range of fragrances and interpretation of this response in consumer terms.

It is well recognised that odours can elicit powerful emotional responses and recall vivid memories. In an attempt better to understand the linkage between odour and emotion researchers have tried to understand the workings of the brain itself. Much has been made of the organisation of the olfactory structures linked directly into the limbic system which itself has been traditionally linked to emotion. However until recently there has been a lack of understanding of brain mechanisms and there has been difficulty in detecting and localising brain activity.

Development of Brain Electrical Activity Mapping (BEAM) by the inventors has involved testing of a wide range of odour types. We have identified significant differences in patterns of brain activity recorded from the surface of the scalp in response to the different odour stimuli. A key question is whether these different patterns of brain activity could give an indication of how people feel in response to fragrance. This was researched by testing a wide range of fragrance types both for patterns of brain activity (using BEAM techniques) and through large scale consumer surveys carried out in the US, France and Germany. The different data sets were compared and correlations were found.

Brain Electrical Activity Mapping

This technique enables measurement of changes in patterns of brain response in real time. The experiment was conducted in a large well ventilated laboratory with the subjects seated comfortably in an armchair. During the experiment subjects experienced reduced visual and auditory perception. To achieve this subjects wore blacked out goggles and headphones which gave a tolerable level of white noise about (70dB). The headphones also allowed the experimenter to communicate with subjects between electroencephalograph (EEG) recording trials. All subjects reported afterwards that they found the experiment to be relaxing and pleasant.

EEG data were collected from over the whole of the scalp using a fabric cap which contained 28 electrodes (Electro-cap International, Inc., USA) built into small plastic buttons in an enhanced International 10/20 electrode placement system. The electrodes were made from highly purified tin and referenced to linked earlobes. A NaCl/KCl gel was used as the conducting medium. The configuration over the scalp is given in Figure 1. Bandpass filters were set at 0.3 and 40 Hz. And sampling rate was 200 Hz. The data handling system dealt with amplitude data from each of the 28 electrodes for all of the conventional EEG wavebands.

A Neuroscience Brain Imager series III model was used to collect the EEG data. This machine uses a computerized real time Fast Fourier Transformation (FFT) technique which allows the averaging of the EEG responses over 2.56 second periods or frames. The EEG signals recorded from the subject are amplified, filtered and then multiplexed prior to analogue to

digital conversion. The Neuroscience Imager gives an averaged EEG amplitude for each electrode shown in Figure 1 every 2.56 s.

EEG data were collected from all of the classical frequency bands (delta, 0-3.5 Hz; theta, 4-7 Hz; alpha 8-13 Hz; beta 1, 15-30 Hz and beta 2 31+ Hz). It was found, in line with earlier similar investigations (e.g. Van Toller et al 1993 (S van Toller, J Behan, P Howells, M Kendal-Reed and A Richardson, "An analysis of spontaneous human cortical EEG activity to odours", *Chemical Senses*, vol. 18, pages 1-16 (1993)), and Moncrieff 1962 (R Moncrieff, "Effects of odours on EEG records", *Perfumery Essential Oil Rec.*, vol. 53, pages 757-760 (1962)) that the predominant change in activity on presentation of the odours was located in the alpha EEG frequency (8-13 Hz). For each electrode a mean amplitude value for the alpha frequencies was calculated across subjects for the first 2.56 second epoch after each stimulus presentation. These means were used to derive a quantitative measure of similarity of EEG responses which were used as input for multivariate analysis eg using Multidimensional Scaling methods (MDS).

Data Analysis

To analyse the data we followed the approach described by Schiffman (see S Schiffman, M Reynolds and F Young, "Introduction to Multidimensional Scaling", Academic Press (1981), and also J Behan and A Richardson, "Sensory Analysis in the Fragrance Industry", *Cosmetics & Toiletries*, 35-39 (1990)). The stimulus space (or map) was derived from a multidimensional scaling (MDS) analysis of the BEAM data. An example of a graphical plot with points for each fragrance given as Figure 2. The contributions of activity at different locations on the scalp was analysed by fitting averaged data for each electrode to the MDS configuration using a vector correlation model (ie the electrode contributions are shown by arrows with the electrode label, O1, FP2, etc, in Figure 2). The same technique was then also used to relate market research data on the fragrance stimuli to the same MDS-derived space (shown as an arrow in the top left hand corner eg 'sensual', 'happy', and in the bottom right hand corner for 'irritating'). By inspection it is possible to deduce the relationship of the various dimensions of response to each other, in particular the relationship between the market research data and the changes in brain wave activity following exposure to perfume.

The MDS map in this case is based on the similarity/dissimilarity of brain response patterns for each perfume. The more different the brain response patterns arising from two of the fragrances, the further apart the two fragrances will appear on the MDS map. The vector model finds the best correlation between the two data sets (ie between the MDS space and the amplitude of response recorded at the electrodes, or between the MDS space and each market research attribute).

Results

Typical BEAM electrode data (in microvolts) are shown in Table 1 for fragrances R2 and R4

The importance of each electrode/attribute to the interpretation of the MDS configuration was determined using formal statistical tests based on F ratios for the vector preference model, as described in the Schiffman reference above. Such an analysis revealed for example that the electrodes Oz, C3, F7, T3, FTC1, F8, T4, FTC2, TCP2 did not meet the set criterion for significance and thus these have not been considered further.

The MDS map was also examined using consumer mood profiling research data, and significant common attributes were identified for the countries where testing was conducted. A summary chart bringing the different facets together is shown in Figure 2 with correlations with German market research data. For the German market research nine of the attributes can be described by the vector model with F-ratios significant at the 5% level. The vectors are shown on the map in Figure 2, pointing in the direction of increasing perception. Maps (not shown) for US and French data were very similar.

Conclusions

By referring to the map shown in Figure 2 and by referring to the related correlation statistics described above we can infer that for this set of fragrances the direction of increasing perception of sensual, happy, warm, calming, safe, reassuring, relaxing and caring across the stimulus space is related to a decrease in activity on electrodes CP1, FP1, C4, F4, TCP1, F3, Fz, and Cz. In view of the wide range of fragrance types studied, the large population of

people included in the consumer research, and similar trends observed in additional studies we have concluded that this is a general trend that can safely be applied to other fragrance sets within a similar odour world.

This and a large amount of other similar data was used to classify fragrance ingredients as R, NR and N, as discussed above, providing the information on which the invention is based.

Table 1: Electrode data		
Electrode	R2	R4
1 (FP1)	13	11
2 (Fz)	14	13
3 (Cz)	14	13
4 (Pz)	20	23
5 (Oz)	14	15
6 (F3)	11	13
7 (C3)	11	10
8 (P3)	17	13
9 (O1)	13	11
10 (F7)	10	10
11 (T3)	8	9
12 (T5)	12	12
13 (FTC1)	11	13
14 (TCP1)	12	12
15 (CP1)	16	14
16 (PO1)	18	15
17 (FP2)	13	12
18 (F4)	14	16
19 (C4)	11	9
20 (P4)	14	13
21 (O2)	10	11
22 (F8)	9	8

23 (T4)	11	8
24 (T6)	13	9
25 (FTC2)	10	9
26 (TCP2)	9	11
27 (CP2)	8	9
28 (PO2)	15	14

CLAIMS

1. A perfume composition comprising:

- a) at least about 25% by weight in total of at least five relaxing fragrance materials (R);
- b) optionally up to about 45% by weight in total of non-relaxing fragrance materials (NR), provided that the weight ratio of R to NR is at least 0.75;
- c) optionally up to about 75% by weight in total of neutral fragrance materials (N);
- d) optionally up to about 25% by weight of other fragrance materials (M) provided that the weight ratio R to (M + NR) is at least 0.75;

and wherein (i) all percentages are based on total weight of the fragrance materials constituting the perfume composition;

(ii) the R materials are selected from the group consisting of anethole, 2-ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)but-2-enol (eg Bangalol™), basil oil, cis-hex-3-enol, coumarin, ethylene brassylate, ethyl linalol, 2-(2'-methylpropyl)-4-hydroxy-4-methyltetrahydropyran (eg Florosa™), hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-benzopyran (eg Galaxolide™), geraniol, cyclohexadecanolide, cyclopentadecanone, methyl anthranilate, alpha-iso-methyl ionone, mixtures of dimethyl benzyl carbinyl butyrate and phenoxyethyl isobutyrate (e.g. Prunella™), mixtures of cyclohexadecanolide and cyclopentadecanone (e.g. Silvanone™), alpha-terpineol, 6-acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin (eg Traseolide™), 2-ethoxy-4-methylphenol (eg Ultravanil™), gamma-undecalactone, vetiver oil, vetiver acetate;

(iii) the NR materials are selected from the group consisting of methyl nonyl aldehyde, allyl amyl glycolate, acetyl cedrene, 3a,6,6,9a-perhydrotetramethylnaphtho [2,1-b] furan (eg Amberlyn Super™), amyl salicylate, armoise oil, benzyl salicylate, bergamot oil, 4-tert-butyl-3-phenylpropanal (eg Bourgeonal™), cedar leaf oil, citronellol, beta-damascone, dimethyl benzyl carbinyl acetate, ethyl 2,6,6-

trimethylcyclohexadienecarboxylate (eg Ethyl Safranate™), methyl 2,4-dihydroxy-3,6-dimethylbenzoate (eg Everniate™), geranyl nitrile, 3-(1,3-benzodioxol-5-yl)-2-methylpropanal (eg Helional™), heliotropin, hexyl salicylate, lemon oil, 2,4-dimethyl-3-cyclohexene-1-carbaldehyde (eg Ligustral™), 3-(4-(1,1-dimethylethyl)phenyl)-2-methylpropanal (eg Lilial™), 4-(4-hydroxy-4-methylpentyl)cyclohex-3-enecarbaldehyde (eg Lyrall™), 3-methyl-5-phenylpentanol (eg Mefrosol™), orange oil, orange terpenes, tagetes oil, tetrahydrogeraniol, vanillin;

(iv) the N materials are selected from the group consisting of benzyl acetate, cassis base, 2-methyl-3-(4-(1-methylethyl)phenyl)propanal (eg Cyclamen Aldehyde™), carvone, cinnamic alcohol, dihydroeugenol, dihydromyrcenol, eugenol, 6-acetyl-1,1,3,4,4,6-hexamethyltetrahydronaphthalene (eg Extralide™), galbanum, gamma-decalactone, hydroxycitronellal, indole, isoeugenol, jasmin oil, 3-pentyltetrahydro-2H-4-pyran ethanoate (eg Jasmopyrane Forte™), linalol, linalyl acetate, methyl dihydrojasmonate, octahydrocoumarin, patchouli oil, 2-phenylethyl alcohol, rose oxide, rose oil, isobornylcyclohexanol (eg Sandalone™), 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol (eg Sandalore™), styrallyl acetate, ylang-ylang;

(v) the M materials comprise perfumery materials not included in the above, excluding odourless or low-odour solvents or diluents used as vehicles for fragrance materials.

2. A perfume composition according to claim 1, which comprises at least 35% by weight in total of relaxing fragrance materials.
3. A perfume composition according to claim 2, which comprises at least 45% by weight in total of relaxing fragrance materials.
4. A perfume composition according to any one of claims 1 to 3, wherein the ratio R to (M+NR) is at least 2.

5. A perfume composition according to any one of the preceding claims, wherein at least 5% by weight of the composition comprises one or more of the following materials: coumarin, ethylene brassylate, hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-benzopyran, cyclohexadecanolide, cyclopentadecanone, 6-acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin, 2-ethoxy-4-methylphenol, gamma-undecalactone.
6. A perfume composition according to any one of the preceding claims, wherein at least 10% by weight of the composition comprises one or more of the following materials: coumarin, ethylene brassylate, hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-benzopyran, cyclohexadecanolide, cyclopentadecanone, 6-acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin, 2-ethoxy-4-methylphenol, gamma-undecalactone.
7. A perfume composition according to any one of the preceding claims, including optionally up to about 35% by weight in total of non-relaxing fragrance materials (NR).
8. A perfume composition according to anyone of the preceding claims, including optionally up to about 25% by weight in total of non-relaxing fragrance materials (NR).
9. A perfume composition according to any one of the preceding claims, wherein the weight ratio of R to NR is at least 0.9 and the weight ratio of R to (M+NR) is at least 0.9.
10. A consumer product comprising a perfume composition in accordance with any one of the preceding claims.
11. A fabric treatment composition comprising a perfume composition in accordance with any one of claims 1 to 9.
12. A personal product composition comprising a perfume composition in accordance with any one of claims 1 to 9.

13. A hard surface treatment composition comprising a perfume composition in accordance with any one of claims 1 to 9.
14. An air care composition comprising a perfume composition in accordance with any one of claims 1 to 9.
15. A method of delivering positive mood benefits particularly relaxation benefits, to a subject, comprising administering to the subject an effective amount of a perfume composition according to any one of claims 1 to 9.
16. A method according to claim 15, wherein the perfume composition is administered in a form selected from the group consisting of solid, liquid, or semisolid compositions intended for application to skin, hair, fabrics or hard surfaces.
17. A method according to claim 15, wherein the perfume composition is administered in a form selected from the group consisting of solid, liquid, or semisolid compositions intended for air freshening.
18. A perfume composition comprising:
- a) from about 25% in total of at least five Relaxing Ingredients (R);
 - b) optionally up to 25% in total of Non-Relaxing Ingredients (NR), provided that the ratio of R to NR is at least 0.9;
 - c) optionally up to 75% in total of Neutral Ingredients (N);
 - d) optionally up to 25% of other perfumery materials (M) provided that the weight ratio R to (M+NR) exceeds 0.9;
- and wherein (i) all percentages are based on weight of the perfume;
- (ii) 'R' ingredients comprise anethole, 2-ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)but-2-enol (eg Bangalol™), basil oil, cis-hex-3-enol, coumarin, ethylene brassylate, ethyl linalol, 2-(2'-methylpropyl)-4-hydroxy-4-methyltetrahydropyran (eg Florosa™), hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]-2-benzopyran (eg Galaxolide™), geraniol, cyclohexadecanolide, cyclopentadecanone, methyl anthranilate,

alpha-iso-methyl ionone, Prunella™, alpha-terpineol, 6-acetyl-1-isopropyl-2,3,3,5-tetrahydrotetralin (eg Traseolide™), 2-ethoxy-4-methylphenol (eg Ultravanil™), gamma-undecalactone, vetiver oil, vetiver acetate;

(iii) 'NR' ingredients comprise methyl nonyl aldehyde, allyl amyl glycolate, acetyl cedrene, 3a,6,6,9a-perhydrotetramethylnaphtho [2,1-b] furan (eg Amberlyn Super™), amyl salicylate, armoise oil, benzyl salicylate, bergamot oil, 4-tert-butyl-3-phenylpropanal (eg Bourgeonal™), cedar leaf oil, citronellol, beta-damascone, dimethyl benzyl carbiny acetate, ethyl 2,6,6-trimethylcyclohexadienecarboxylate (eg Ethyl Safranate™), methyl 2,4-dihydroxy-3,6-dimethylbenzoate (eg Everniate™), geranyl nitrile, 3-(1,3-benzodioxol-5-yl)-2-methylpropanal (eg Helional™), heliotropin, hexyl salicylate, lemon oil, 2,4-dimethyl-3-cyclohexene-1-carbaldehyde (eg Ligustral™), 3-(4-(1,1-dimethylethyl)phenyl)-2-methylpropanal (eg Lilial™), 4-(4-hydroxy-4-methylpentyl)cyclohex-3-enecarbaldehyde (eg Lyrall™), 3-methyl-5-phenylpentanol (eg Mefrosol™), orange oil, tagetes oil, tetrahydrogeraniol, vanillin;

(iv) 'N' ingredients comprise benzyl acetate, cassis base, 2-methyl-3-(4-(1-methylethyl)phenyl)propanal (eg Cyclamen Aldehyde™), carvone, dihydroeugenol, dihydromyrcenol, eugenol, 6-acetyl-1,1,3,4,4,6-hexamethyltetrahydronaphthalene (eg Extralide™), gamma-decalactone, hydroxycitronellal, indole, isoeugenol, jasmin oil, 3-pentyltetrahydro-2H-4-pyranyl ethanoate (eg Jasmopyrane Forte™), linalol, linalyl acetate, methyl dihydrojasmonate, octahydrocoumarin, patchouli oil, 2-phenylethyl alcohol, rose oxide, rose oil, isobornylcyclohexanol (eg Sandalone™), 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol (eg Sandalore™), styrallyl acetate, ylang-ylang;

(v) 'M' ingredients comprise perfumery materials not included in the above, excluding odourless or low-odour solvents or diluents used as vehicles for other perfume components.

Fig.1.





Fig.2.



INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB 01/05589

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K7/46 C11B9/00 A61K7/06 A61K7/48 A61L9/00
C11D3/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C11B A61L C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	US 6 268 333 B1 (NAKAMURA SHOJI ET AL) 31 July 2001 (2001-07-31) the whole document	1-18
A	& JP 06 172781 A 21 June 1994 (1994-06-21) cited in the application	
Y	DE 41 25 562 A (EBERT KARIN) 10 December 1992 (1992-12-10) abstract column 3, line 6 - line 22	1-18
A	EP 0 978 273 A (SHISEIDO CO LTD) 9 February 2000 (2000-02-09) page 3, line 32 - line 48	1-18
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

8 February 2002

Date of mailing of the international search report

22/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Dauksch, H

INTERNATIONAL SEARCH REPORT

Int Application No
PCT/GB 01/05589

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE WPI Section Ch, Week 199806 Derwent Publications Ltd., London, GB; Class D21, AN 1998-059489 XP002189736 & JP 09 302377 A (POLA CHEM IND INC), 25 November 1997 (1997-11-25) cited in the application abstract ---	1-18
A	DATABASE WPI Section Ch, Week 198839 Derwent Publications Ltd., London, GB; Class D21, AN 1988-274031 XP002189737 & JP 63 199292 A (TAKASAGO PERFUMERY CO LTD), 17 August 1988 (1988-08-17) abstract ---	1-18
A	US 5 904 916 A (HIRSCH ALAN R) 18 May 1999 (1999-05-18) cited in the application abstract ---	1
A	PATENT ABSTRACTS OF JAPAN vol. 1998, no. 01, 30 January 1998 (1998-01-30) & JP 09 227399 A (POLA CHEM IND INC), 2 September 1997 (1997-09-02) cited in the application abstract -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB 01/05589

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6268333	B1	31-07-2001	JP 6172781 A	21-06-1994
DE 4125562	A	10-12-1992	DE 4125562 A1	10-12-1992
EP 0978273	A	09-02-2000	JP 2000053532 A	22-02-2000
			CN 1280859 A	24-01-2001
			EP 0978273 A1	09-02-2000
JP 9302377	A	25-11-1997	NONE	
JP 63199292	A	17-08-1988	NONE	
US 5904916	A	18-05-1999	NONE	
JP 09227399	A	02-09-1997	NONE	